

Do Not Use Abbreviations: U (for Unit), IU for International unit), Q.D., Q.O.D., Trailing Zero (X.0 mg) MS, MSO4 MgSO4

PRE-OPERATIVE ORDERS

- Status: Admit to Inpatient Status (I certify that inpatient services are needed)
 Place Patient in Outpatient Status
 Place Patient in Outpatient Status and begin Observation Services

Admit to the service of:

PATIENT NAME (LAST):		FIRST NAME	DATE OF BIRTH:
DIAGNOSIS:			ANESTHESIA TYPE:
PROCEDURE CONSENT TO STATE:			
DATE OF SURGERY/PROCEDURE	PHYSICIAN:	PRIMARY PHYSICIAN:	CPT CODES:
CPT CODE(S)			

ALLERGIE(S) Type of Reaction(s): Patient Weight: _____ kg
--

IV FLUIDS:

- Lactated Ringers @ 30 mL/hr on arrival to Preop
 0.9% Sodium Chloride @ 30 mL/hr on arrival to Preop

Preop antibiotics:

- Cefazolin 1 gm IV for patient weight < 60 kg, infuse within 60 minutes prior to surgery
 Cefazolin 2 gm IV for patient weight 60-120 kg, infuse within 60 minutes prior to surgery
 Cefazolin 3 gm IV for patient weight > 120 kg, infuse within 60 minutes prior to surgery

If beta-lactam allergy or has a history or risk for MRSA, instead of cefazolin, give

- Vancomycin 750 mg for patient weight < 50 kg IV over 60 minutes, infuse within 120 minutes prior to surgery
 Vancomycin 1 gm for patient weight 50 - 100 kg IV over 60 minutes, infuse within 120 minutes prior to surgery
 Vancomycin 1.5 gm for patient weight > 100 kg IV over 90 minutes, infuse within 120 minutes prior to surgery

If beta-lactam and vancomycin intolerant, give clindamycin instead of cefazolin or vancomycin:

- Clindamycin 900 mg IV over 30 minutes, start 60 minutes prior to surgery

Cardiac or Vascular Surgery:

- Cefazolin dose as above x 1 preop
 Vancomycin dose as above x 1 preop
 Clindamycin dose as above x 1 preop Intra-abdominal Surgery:
 Cefazolin, dose as above and metronidazole 500 mg IV x 1 dose each preop
 Levofloxacin 500 mg IV and metronidazole 500 mg IV x 1 dose each preop

Gynecologic Surgery:

- Cefazolin, dose as above x 1 preop
 Clindamycin, dose as above x 1 preop
 Vancomycin, dose as above x 1 preop
 Other medication order: _____

Medications:

A. To be given in pre-op day of procedure

B. Patient given prescription to take the medication prior to arrival for surgery

- Acetaminophen 975 mg PO x 1
 Acetaminophen 650 mg liquid PO x 1
 Acetaminophen 1gm IV x 1
 Celecoxib 200 mg PO x 1 preop
 Gabapentin (Neurotin) 600 mg PO x 1
 Gabapentin (Neurotin) 200 mg PO x 1
Reminder: If age > 75, patient on dialysis, or <50kg weight, give:
 Gabapentin (Neurotin) 300 mg PO x 1 preop
 Oxycodone SUSTAINED release (Oxycontin) 10 mg PO x 1
 Oxycodone IMMEDIATE release (OxyIR) 10 mg PO x 1
 Metoclopramide 10 mg IV x 1 dose
 Other medication order: _____

- Tramadol 50mg PO x 1
 Dexamethasone 8mg x 1 (DO NOT ORDER IF DIABETIC)

Physician Signature: _____

Date/Time: ____ / ____ / ____ at: _____

PRE OPERATIVE ORDERS



5301 South Congress Avenue
Atlantis, FL 33462

Patient Identification/Label

PRE-OPERATIVE ORDERS

EKG Done at: JFK PCP **Must Be Legible Copy**

Labs Done at: JFK
 Outside Testing

Please use Anesthesia Guidelines to determine testing.

- A1C
- CBC CBC w/Differential
- Chem 7 PT, PTT & INR
- Chem 25 Liver Profile
- Sickle Cell BHCG < 55 yrs.

- Urinalysis CEA
- Urine Culture & Sensitivity
- Type & Screen
- MRSA/MSSA Screening
- Type & Cross X _____ units

PTH Analyzer:

- Hematology Testing
- Nuclear Medicine Injection

Other Labs: _____

- Anti Embolic Hose
- Sequential Compression Device(s) Case Management to Arrange: _____

- Incentive Spirometer

Medical Pre Op Evaluation: Phone: _____
 No Yes Dr.:

Cardiac Pre Op Evaluation: Phone: _____
 No Yes Dr.:

Other Pre Op Evaluation (Type): Phone: _____
 No Yes Dr.:

Patient From Nursing Home/Extended Care Facility? No Yes Phone: _____
 Name: _____

NPO AFTER MIDNIGHT, DATE:
 ENHANCED SURGICAL RECOVERY

Diet:

- No solid food after midnight the night before the procedure unless otherwise instructed by anesthesia.
- May have clear liquids (NO RED COLOR OR DYE) up to arrival time at JFK or until 2 hours before scheduled surgery.
- If instructed to do bowel prep prior to surgery, no solid food starting at midnight 2 nights prior to surgery.
- INSTRUCT PATIENT TO DRINK pre-surgery drink:**
 - Drink 2 bottles evening prior to surgery and drink one bottle at least 2 hours prior to scheduled surgery time.
 - If patient is Diabetic**, substitute Gatorade Zero for pre-surgery drink and instruct to drink one 20 oz. bottle the evening prior to procedure and one-half bottle of Gatorade zero 2 hours prior to scheduled procedure.
- Instruct patient to shower/bathe with 2% chlorhexidine gluconate (CHG) shower soap the night before surgery and repeat the morning of surgery.
- Upon arrival to preop have patient wipe body down with 2% chlorhexidine gluconate (CHG) wipes.

Chest X-Ray
 JFK Outside testing

KUB day of procedure: _____
 Breast: _____
 MRI: _____
 CT: _____

Obtain Test Results:

MRA VEIN MAPPING OTHER: _____

DONE AT: _____

Other: _____

PERSON COMPLETING FORM: NAME (PLEASE PRINT): _____ DATE: _____ TIME: _____

PHYSICIAN'S SIGNATURE: PHYSICIAN'S NAME (PLEASE PRINT): _____ DATE: _____ TIME: _____

PRE OPERATIVE ORDERS



Patient Identification/Label